



E2088

JACC March 12, 2013

Volume 61, Issue 10



Vascular Medicine

AMNIOTIC MESENCHYMAL STEM CELLS WITH ROBUST CHEMOTACTIC PROPERTIES ARE EFFECTIVE IN THE TREATMENT OF A MYOCARDIAL INFARCTION MODEL

Poster Contributions

Poster Sessions, Expo North

Sunday, March 10, 2013, 3:45 p.m.-4:30 p.m.

Session Title: Angiogenesis and Vascular Injury

Abstract Category: 33. Vascular Medicine: Basic

Presentation Number: 1254-170

Authors: Sung-Whan Kim, Hong-Zhe Zhang, Kim Moo Hyun, Dong-A University Hospital, Busan, South Korea

Background: We previously reported that amniotic mesenchymal stem cells (AMMs) possess high angio-vasulogenic properties. In this study, we investigated the chemotactic abilities of AMMs for improved cardiac function and regenerative angiogenesis.

Methods: The expressions of chemotactic and angiogenic genes were determined by qRT-PCR. Myocardial infarction (MI) was induced in NOD/SCID mice and cells were directly transplanted into the border regions of ischemic heart tissue. Immunohistochemical analysis was also conducted.

Results: AMMs significantly expressed the representative chemotactic factor GCP-2, NAP-2 as well as angiogenic factor Hif-1 α . AMMs also highly expressed the chemokine receptors CCR2, CCR3 and CCR5. AMM transplantation improved left ventricular function, capillary density, angiogenic cytokine levels, angiopoietin (Ang)-1 and vascular endothelial growth factor (VEGF-A) levels in affected tissue. Immunohistochemical assaying also revealed increased engraftment and endothelial phenotypes.

Conclusions: Our findings suggest that due to elevated survival and related chemotactic potential, AMMs are a promising stem cell source for the treatment of ischemic cardiovascular disease.